

# Calculus of Variations

## Summer Term 2017

### Lecture 3

Universität des Saarlandes

28 April 2017

## Purpose of Lesson:

- To continue the discussion about special cases of the E-L equation.
- To show that the E-L equation is a necessary, but not sufficient condition for a local extremum.
- To discuss the simplest variational problems involving undetermined end points.

## Special Cases of the E-L Equation (cont.)

3. If  $F$  does not depend on  $y'$ , the E-L equation takes the form

$$F_y(x, y) = 0,$$

and hence is not a differential equation, but a *finite*, whose solution consists of one or more curves  $y = y(x)$

4. In a variety of problems, one encounters functionals of the form

$$J[y] = \int_a^b f(x, y) \sqrt{1 + y'^2} dx,$$

representing the integral of a function  $f(x, y)$  with respect to the **arc length**  $s$  ( $ds = \sqrt{1 + y'^2} dx$ ).

- In this case, the E-L equation can be transformed into

$$\begin{aligned} F_y - \frac{d}{dx} F_{y'} &= f_y(x, y) \sqrt{1 + y'^2} - \frac{d}{dx} \left[ f(x, y) \frac{y'}{\sqrt{1 + y'^2}} \right] \\ &= \frac{1}{\sqrt{1 + y'^2}} \left[ f_y - f_x y' - f \frac{y''}{1 + y'^2} \right] = 0 \end{aligned}$$

i.e.,


$$f_y - f_x y' - f \frac{y''}{1 + y'^2} = 0.$$

# Failure of Sufficiency

## Remark

Note that the Euler-Lagrange Equation is a **necessary**, but not **sufficient** condition for a **local** extremum.

Next we will consider the famous example of the minimal surface area for a soap film.



## Example 2.1

We want to minimize the following functional

$$J[y] = 2\pi \int_{x_0}^{x_1} y \sqrt{1 + (y')^2} dx \rightarrow \min$$

according to the boundary conditions  $y(x_0) = y_0$ ,  $y(x_1) = y_1$ .

- If we use the Euler-Lagrange equation and solve it for  $y(x)$  we find the **Catenary** function

$$y(x) = C_1 \cosh \left\{ \left( \frac{x + C_2}{C_1} \right) \right\}$$

- Consider the special case where  $C_2 = 0$  and require that  $y(x) = C_1 \cosh \left\{ \left( \frac{x}{C_1} \right) \right\}$  pass through  $(-x_1, 1)$  and  $(x_1, 1)$  where  $x_1$  is a constant.
- So  $C_1$  will satisfy

$$1 = C_1 \cosh \left\{ \left( \frac{x}{C_1} \right) \right\} \quad (2.3)$$

- Compare  $y = 1$  and (2.3) versus  $C_1$ 
  - 1 For  $x_1 = 1$  there is **No Solutions**;
  - 2 For  $x_1 = 0.7$  there is exactly **One Solutions**;
  - 3 For  $x_1 = 0.4$  there are **Two Solutions**.

### §3. Undetermined End Points



# Undetermined End Points

## Problem 3-1

We seek to minimize the integral

$$J[y] = \int_a^b F(x, y, y') dx \rightarrow \min$$

with respect to functions that attain the value  $A$  for  $x = a$ , but for which no value is prescribed at  $x = b$ .

## Question:

What is the arc of quickest descent from a fixed point to a vertical line?

- To find the minimizing function, we as before introduce a small variation from  $y(x)$ , namely

$$y(x) + \varepsilon\eta(x)$$

where  $\varepsilon$  is a small parameter and  $\eta(x)$  is a smooth curve satisfying the BC

$$\eta(a) = 0.$$

- We take the derivative of

$$\phi(\varepsilon) = J[y + \varepsilon\eta]$$

with respect to  $\varepsilon$ , evaluate it  $\varepsilon = 0$ , and set this equal to zero; that is,

$$\begin{aligned} \frac{d\phi(\varepsilon)}{d\varepsilon} &= \frac{d}{d\varepsilon} J[y + \varepsilon\eta] \Big|_{\varepsilon=0} \\ &= \int_a^b \left[ \frac{\partial F}{\partial y} \eta(x) + \frac{\partial F}{\partial y'} \eta'(x) \right] dx \end{aligned}$$

- Integration by parts gives

$$\left[ \frac{\partial F}{\partial y'} \eta(x) \right]_{x=a}^{x=b} + \int_a^b \left\{ \frac{\partial F}{\partial y} - \frac{d}{dx} \left[ \frac{\partial F}{\partial y'} \right] \right\} \eta(x) dx = 0. \quad (3.1)$$

- Since (3.1) must hold for **all** choices of  $\eta(x)$  satisfying

$$\eta(a) = 0,$$

it must in particular hold for **those**  $\eta$  for which  $\eta(b) = 0$ . For such  $\eta(x)$  the first term in (3.1) disappeared and as before we end up with

$$\boxed{\frac{\partial F}{\partial y} - \frac{d}{dx} \left[ \frac{\partial F}{\partial y'} \right] = 0}. \quad (3.2)$$

- So, with result (3.2), and for **general**  $\eta(x)$  once again, the second member of (3.1) reduces to its first term

$$\left[ \frac{\partial F}{\partial y'} \eta(x) \right]^{x=b} = 0.$$

- Now, by choosing  $\eta(b) = 1$ , the vanishing for all  $\eta$  of the term remaining requires fulfillment of the **end-point condition**

$$\boxed{\left. \frac{\partial F}{\partial y'} \right|_{x=b} = 0}. \quad (3.3)$$

- The two constants of integration obtained in the solution of (3.2), a second-order equation, are determined by the end-point condition  $y(a) = A$  and (3.3) - provided, of course, a solution of the problem exists.

## Problem 3-2

We seek to minimize the integral

$$J[y] = \int_a^{x^*} F(x, y, y') dx \rightarrow \min$$

with respect to functions which attain the value  $A$  for  $x = a$  and which satisfy the given relation

$$g(x, y) = 0$$

at the upper limit of integration, as yet undetermined.

### Question:

- What is the arc of quickest descent from a fixed point to a given curve?

- To find the minimizing function, we as before introduce a small variation from  $y(x)$ , namely

$$y(x) + \varepsilon\eta(x)$$

where  $\varepsilon$  is a small parameter and  $\eta(x)$  is a smooth curve satisfying the BC

$$\eta(a) = 0.$$

- The point of intersection of our small variation  $y(x) + \varepsilon\eta(x)$  with the given curve  $g(x, y) = 0$  is denoted by  $(x^*, y^*)$ . We thus have

$$\boxed{g(x^*, y^*) = 0, \quad y^* = y(x^*) + \varepsilon\eta(x^*)}. \quad (3.4)$$

- We take the derivative of

$$\phi(\varepsilon) = J[y + \varepsilon\eta]$$

with respect to  $\varepsilon$ , evaluate it  $\varepsilon = 0$ , and set this equal to zero; that is,

$$\begin{aligned} \frac{d\phi(\varepsilon)}{d\varepsilon} &= \frac{d}{d\varepsilon} J[y + \varepsilon\eta] \Big|_{\varepsilon=0} \\ &= \int_a^{x^*} \left[ \frac{\partial F}{\partial y} \eta(x) + \frac{\partial F}{\partial y'} \eta'(x) \right] dx + F(x^*, y^*, (y^*)') \frac{dx^*}{d\varepsilon} \Big|_{\varepsilon=0} \end{aligned}$$

- Since relations (3.4) hold for **all**  $\varepsilon$ , we have that the total derivative of  $g(x^*, y^*)$  with respect to  $\varepsilon$  must vanish.
- From (3.4) we therefore obtain, on noting that  $x^*$  is a function of  $\varepsilon$  for any given  $\eta(x)$ ,

$$\begin{aligned} 0 &= \frac{\partial g}{\partial x^*} \frac{dx^*}{d\varepsilon} + \frac{\partial g}{\partial y^*} \frac{dy^*}{d\varepsilon} \\ &= \frac{\partial g}{\partial x^*} \frac{dx^*}{d\varepsilon} + \frac{\partial g}{\partial y^*} \left[ y'(x^*) \frac{dx^*}{d\varepsilon} + \eta(x^*) + \varepsilon \eta'(x^*) \frac{dx^*}{d\varepsilon} \right]. \end{aligned}$$

- Solving the above equality, with  $\varepsilon = 0$ , for  $\left. \frac{dx^*}{d\varepsilon} \right|_{\varepsilon=0}$  we obtain

$$\left. \frac{dx^*}{d\varepsilon} \right|_{\varepsilon=0} = - \frac{\eta(x^*) \frac{\partial g}{\partial y^*}}{\frac{\partial g}{\partial x^*} + y'(x^*) \frac{\partial g}{\partial y^*}} \quad (3.5)$$



- With the aid of (3.5) and integration by parts we get

$$\int_a^{x^*} \left[ \frac{\partial F}{\partial y} - \frac{d}{dx} \left( \frac{\partial F}{\partial y'} \right) \right] \eta dx + \eta(x^*) \left[ \frac{\partial F}{\partial y'} - \frac{F \frac{\partial g}{\partial y^*}}{\frac{\partial g}{\partial x^*} + (y^*)' \frac{\partial g}{\partial y^*}} \right]^{x=x^*} = 0$$

- Repeating the line of arguments carried out in Problem 3-1 above we conclude that  $y = y(x)$  satisfies the Euler-Lagrange equation

$$\frac{\partial F}{\partial y} - \frac{d}{dx} \left[ \frac{\partial F}{\partial y'} \right] = 0$$

and, in addition to the BC  $y(a) = A$ , the right-hand end-point condition

$$\left[ \frac{\partial F}{\partial y'} - \frac{F \frac{\partial g}{\partial y^*}}{\frac{\partial g}{\partial x^*} + (y^*)' \frac{\partial g}{\partial y^*}} \right]^{x=x^*} = 0.$$