



Calculus of Variations (Summer Term 2016)
Assignment H4 - Homework

Problem 4.1 (8 Points)

Minimize the functional

$$J[y] = \int_0^2 (y' + 1)^2 (y')^2 dx$$

subject to the end-point conditions that $x(0) = 1$ and $x(2) = 0$.

(*Hint*: consider the possibility of broken extremals).

Problem 4.2 (3x3=9 Points)

Decide which of the following functionals admit broken extremals or not:

$$J_1[y] = \int_0^1 (y'^6 - 5y'^4 + 15y'^2) dx, \quad J_2[y] = \int_0^1 y'^3 dx,$$

$$J_3[y] = \int_0^1 \cos \{(y')\} dx.$$

(Either construct examples of broken extremals or rule out their existence using available theory). Moreover, determine at least one admissible extremal for the minimization of each of the functionals over the set of functions y satisfying $y(0) = 0$ and $y(1) = \pi/2$.

Problem 4.3 (8 Points)

Use Ritz's method to find an approximate solution to minimize the

$$J[y] = \int_0^{2\pi} (y'^2 + \lambda^2 y^2) dx, \quad y(0) = 0, \quad y(1) = 1$$

where $y(0) = 1$ and $y(2\pi) = 1$ and λ is a positive integer. Use the trial functions

$$\phi_n(x) = \cos(nx).$$

Compare your solution to one found directly from the Euler-Lagrange equations.

Deadline for submission: Tuesday, June 21, 08:30 am