

# PDE and Boundary-Value Problems

## Winter Term 2016/2017

### Lecture 18

Saarland University

03 February 2017

## Purpose of Lesson

- To solve the Dirichlet problem between two circles (annulus).
- To discuss briefly the solution to the exterior Dirichlet problem for the circle.
- To find particular solutions of the Laplace equation in spherical coordinates. To solve the interior and exterior Dirichlet problems for the Laplace equation in 3D.

# The Dirichlet Problem in an Annulus

## Problem 18-1

To find the function  $u(r, \theta)$  that satisfies

$$\text{PDE: } u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0, \quad R_1 < r < R_2$$

$$\text{BCs: } \begin{cases} u(R_1, \theta) = g_1(\theta), \\ u(R_2, \theta) = g_2(\theta) \end{cases} \quad 0 \leq \theta < 2\pi.$$

## Step 1. (Separation of Variables)

- Substituting  $u(r, \theta) = R(r)\Theta(\theta)$  into the Laplace equation and arguing similar to the interior Dirichlet problem we arrive at our general solution

$$\begin{aligned} u(r, \theta) = & a_0 + b_0 \ln r \\ & + \sum_{n=1}^{\infty} (a_n r^n + b_n r^{-n}) \cos(n\theta) \\ & + \sum_{n=1}^{\infty} (c_n r^n + d_n r^{-n}) \sin(n\theta) \end{aligned} \tag{18.1}$$

## Step 2. (Substituting into BCs)

- Substituting the solution (18.1) into the BCs and integrating gives the following equations:

$$\left\{ \begin{array}{l} a_0 + b_0 \ln R_1 = \frac{1}{2\pi} \int_0^{2\pi} g_1(s) ds \\ a_0 + b_0 \ln R_2 = \frac{1}{2\pi} \int_0^{2\pi} g_2(s) ds \end{array} \right. \quad (\text{Solve for } a_0, b_0)$$

## Step 2. (cont.)

$$\left\{ \begin{array}{l} a_n R_1^n + b_n R_1^{-n} = \frac{1}{\pi} \int_0^{2\pi} g_1(s) \cos(ns) ds \\ a_n R_2^n + b_n R_2^{-n} = \frac{1}{\pi} \int_0^{2\pi} g_2(s) \cos(ns) ds \end{array} \right. \quad (\text{Solve for } a_n, b_n)$$

$$\left\{ \begin{array}{l} c_n R_1^n + d_n R_1^{-n} = \frac{1}{\pi} \int_0^{2\pi} g_1(s) \sin(ns) ds \\ c_n R_2^n + d_n R_2^{-n} = \frac{1}{\pi} \int_0^{2\pi} g_2(s) \sin(ns) ds \end{array} \right. \quad (\text{Solve for } c_n, d_n)$$

# Exterior Dirichlet Problem

## Problem 18-2

To find the function  $u(r, \theta)$  that satisfies

$$\text{PDE: } u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0, \quad 1 < r < \infty$$

$$\text{BC: } u(1, \theta) = g(\theta), \quad 0 \leq \theta < 2\pi.$$

Problem 18-2 is solved exactly like the interior Dirichlet problem. The only exception is that now we throw out the solutions that are **unbounded** as  $r$  goes to **infinity**

$$r^n \cos(n\theta), \quad r^n \sin(n\theta), \quad \ln r$$

# Exterior Dirichlet Problem (cont.)

Hence, we are left with the solution

$$u(r, \theta) = \sum_{n=0}^{\infty} r^{-n} [a_n \cos(n\theta) + b_n \sin(n\theta)],$$

where  $a_n$  and  $b_n$  are exactly as before

$$a_0 = \frac{1}{2\pi} \int_0^{2\pi} g(s) ds,$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} g(s) \cos(ns) ds, \quad b_n = \frac{1}{\pi} \int_0^{2\pi} g(s) \sin(ns) ds$$



## Remarks

- The exterior Dirichlet problem for arbitrary radius  $R$

$$\text{PDE: } u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0, \quad R < r < \infty$$

$$\text{BC: } u(R, \theta) = g(\theta), \quad 0 \leq \theta < 2\pi.$$

has the solution

$$u(r, \theta) = \sum_{n=0}^{\infty} \left(\frac{r}{R}\right)^{-n} [a_n \cos(n\theta) + b_n \sin(n\theta)]$$

- The only solution of the 2-D Laplace equation that depend only on  $r$  are **constants** and  **$\ln r$** . The potential  $\ln r$  is very important and is called the **logarithmic potential**.

# Laplace's Equation in Spherical Coordinates (Spherical Harmonics)

An important problem is to find the potential inside or outside a sphere when the potential is given on the boundary. Consider, first, the **interior problem**:

## Problem 18-3

To find the function  $u(r, \theta, \phi)$  that satisfies

$$\text{PDE: } (r^2 u_r)_r + \frac{1}{\sin \phi} [\sin \phi u_\phi]_\phi + \frac{1}{\sin^2 \phi} u_{\theta\theta} = 0, \quad 0 < r < 1$$

$$\text{BC: } u(1, \theta, \phi) = g(\theta, \phi), \quad -\pi \leq \theta < \pi, \quad 0 \leq \phi < \pi$$

## Remarks

- A typical application of the problem 18-3 would be to find the temperature inside a sphere when the temperature is specified on the boundary.
- Quite often  $g(\theta, \phi)$  has a **specific form**, so that it isn't necessary to solve the problem in its most general form.
- We consider two important cases. One is the case when  $g(\theta, \phi)$  is **constant**, and the other is when it depends **only** on the angle  $\phi$  (the angle from the north pole).

## Special Case 1. ( $g(\theta, \phi) = \text{constant}$ )

- In this case, it is clear that the solution is independent of  $\theta$  and  $\phi$ , and so Laplace's equation reduces to the ODE

$$(r^2 u_r)_r = 0. \quad (18.2)$$

- The general solution of (18.2) is

$$u(r) = \frac{a}{r} + b$$

- In other words, constants and  $\frac{c}{r}$  are the only potentials that depend only on the radial distance from the origin. The potential  $\frac{1}{r}$  is called the **Newtonian potential**.

## Special Case 2. ( $g(\theta, \phi)$ depends only on $\phi$ )

In this case, the Dirichlet problem takes the form

### Problem 18-3a

To find the function  $u(r, \theta, \phi)$  that satisfies

$$\text{PDE: } (r^2 u_r)_r + \frac{1}{\sin \phi} [\sin \phi u_\phi]_\phi = 0, \quad 0 < r < 1$$

$$\text{BC: } u(1, \theta, \phi) = g(\phi), \quad 0 \leq \phi < \pi$$

## Step 1. (Separation of variables)

- We look for solutions of the form

$$u(r, \phi) = R(r)\Phi(\phi)$$

and arrive at the two ODEs

$$r^2 R'' + 2rR' - n(n+1)R = 0 \quad (\text{Euler's equation})$$

$$[\sin \phi \Phi']' + n(n+1) \sin \phi \Phi = 0 \quad (\text{Legendre's equation})$$

- The separation constant is chosen to be  $-n(n+1)$  for convenience; later we will see why this choice is made.

## Step 2. (Solving the Euler equation)

- We solve Euler's equation by substituting  $R(r) = r^\alpha$  in the equation and solving for  $\alpha$ . Doing this, we get two values

$$\alpha = \begin{cases} n \\ -(n+1) \end{cases}$$

- Hence, Euler's equation has the general solution

$$R(r) = ar^n + br^{-(n+1)}$$

### Step 3. (Solving the Legendre equation)

- Making the substitution  $x = \cos \phi$  we get the new Legendre equation

$$(1 - x^2) \frac{d^2 \Phi}{dx^2} - 2x \frac{d\Phi}{dx} + n(n + 1)\Phi = 0, \quad -1 \leq x \leq 1.$$

The idea here is to solve for  $\Phi(x)$  and then substitute  $x = \cos \phi$  in the solution.

- Legendre's equation is a linear second-order ODE with variable coefficients. One of the difficulties in this equation is that the coefficient  $(1 - x^2)$  is zero at the ends of the interval  $[-1, 1]$ . Equations like this are called **singular differential equations** and are often solved by the **method of Frobenius**.



## Step 3. (cont.)

- The only bounded solutions of Legendre's equation occur when  $n = 0, 1, 2, \dots$  and these solutions are **polynomials**  $P_n(x)$ ,  $-1 \leq x \leq 1$  (Legendre polynomials)

$$n = 0 \quad P_0(x) = 1$$


$$n = 1 \quad P_1(x) = x$$

$$n = 2 \quad P_2(x) = \frac{1}{2}(3x^2 - 1)$$

$$n = 3 \quad P_3(x) = \frac{1}{2}(5x^2 - 3x)$$

$$\vdots \quad \vdots \quad \quad \quad \vdots$$

$$n \quad P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n] \quad (\text{Rodrigues' formula})$$

Legendre Polynomials  $P_n(x)$ 

## Step 4. (Combination)

- We now have that the bounded solutions of

$$r^2 R'' + 2rR' - n(n+1)R = 0 \quad 0 < r < 1$$

$$[\sin \phi \Phi']' + n(n+1) \sin \phi \Phi = 0 \quad -\pi \leq \phi \leq \pi$$

are

$$R(r) = ar^n$$

$$\Phi(\phi) = aP_n(\cos \phi)$$

- Therefore,

$$u(r, \phi) = \sum_{n=0}^{\infty} a_n r^n P_n(\cos \phi). \quad (18.3)$$

## Step 5. (Substituting into BC)

- Substituting solution (18.3) into the BC gives

$$\sum_{n=0}^{\infty} a_n P_n(\cos \phi) = g(\phi) \quad (18.4)$$

- Observe that the Legendre polynomials are orthogonal on  $[-1, 1]$ .

## Step 5. (cont.)

So, if we multiply each side of (18.4) by  $P_m(\cos \phi) \sin \phi$  and integrate  $\phi$  from 0 to  $\pi$ , we get

$$\begin{aligned} \int_0^\pi g(\phi) P_m(\cos \phi) \sin \phi d\phi &= \sum_{n=0}^{\infty} a_n \int_0^\pi P_n(\cos \phi) P_m(\cos \phi) \sin \phi d\phi \\ &= \sum_{n=0}^{\infty} a_n \int_{-1}^1 P_n(x) P_m(x) dx \\ &= \begin{cases} 0, & n \neq m \\ \frac{2a_m}{2m+1}, & m = n \end{cases} \end{aligned}$$

## Step 5. (cont.)

- Hence

$$a_n = \frac{2n+1}{2} \int_0^\pi g(\phi) P_n(\cos \phi) \sin \phi d\phi$$

and the solution to our Dirichlet problem 18-3a is

$$u(r, \phi) = \sum_{n=0}^{\infty} a_n r^n P_n(\cos \phi)$$

## Remarks

- The solution of the exterior Dirichlet problem

$$\text{PDE: } \Delta u = 0, \quad 1 < r < \infty$$

$$\text{BC: } u(1, \theta, \phi) = g(\phi), \quad 0 \leq \phi < \pi$$

is

$$u(r, \phi) = \sum_{n=0}^{\infty} \frac{b_n}{r^{n+1}} P_n(\cos \phi),$$

where

$$b_n = \frac{2n+1}{2} \int_0^{\pi} g(\phi) P_n(\cos \phi) \sin \phi d\phi.$$

## Remarks (cont.)

- For example, the BC  $g(\phi) = 3$  would yield for the solution of the exterior problem

$$u(r, \phi) = \frac{3}{r}.$$

Note that in this problem (in 3D!!!), the solution goes to zero, while in **two dimensions**, the exterior solution with constant BC was **itself** a constant.